# **VARDENAFIL (Levitra) Fact Sheet [G]**

### **Bottom Line:**

Vardenafil doesn't offer any benefits compared to sildenafil, which has more data in the psychiatric setting and more clinical experience.

#### **FDA Indications:**

**Erectile dysfunction.** 

# **Dosage Forms:**

- Tablets (Levitra, [G]): 2.5 mg, 5 mg, 10 mg, 20 mg.
- Orally disintegrating tablets (Staxyn, [G]): 10 mg.

# **Dosage Guidance:**

- Start 10 mg ×1 one hour prior to sexual activity (5 mg if >65 years old). Max 20 mg/dose and one dose/day.
- Orally disintegrating tablets (Staxyn): Start 10 mg ×1 one hour prior to sexual activity. Max 10 mg/day.
- Taking with a high-fat meal may decrease serum vardenafil levels by as much as 50%.

Monitoring: No routine monitoring recommended unless clinical picture warrants.

### Cost: \$

## **Side Effects:**

- Most common: Flushing, headache, nasal congestion, heartburn.
- Serious but rare: May cause dose-related impairment of color discrimination. Sudden decrease or loss of hearing has been reported rarely; hearing changes may be accompanied by tinnitus and dizziness. Decreases in blood pressure may occur due to vasodilator effects; concurrent use with alpha-adrenergic antagonists or substantial alcohol consumption may cause symptomatic hypotension. Avoid use with nitrates (see below). Painful erection greater than six hours in duration (priapism) may occur rarely.

# Mechanism, Pharmacokinetics, and Drug Interactions:

- Phosphodiesterase type 5 (PDE-5) inhibitor.
- Metabolized primarily through CYP3A4 and to a lesser extent 2C9; t ½: 4–5 hours.
- Avoid concomitant use with nitrates in any form (eg, nitroglycerin, isosorbide dinitrate, amyl nitrite "poppers"). Use with caution in patients taking alpha blockers; may cause symptomatic hypotension (use 5 mg dose). Use with caution in patients taking strong CYP3A4 inhibitors, which may increase or extend effects of vardenafil (maximum of 2.5 mg/day).

# **Clinical Pearl:**

Usual onset of effect of vardenafil is within one hour of a dose; effects usually last two hours.

Staxyn, the orally disintegrating tablet form of vardenafil, is peppermint flavored.

